arXiv:1510.08168v1 [math.CO] 28 Oct 2015

On two q-ary n-cube coloring problems

Zhe Han* **Mei Lu†**
Department of Mathematical Sciences, Tsinghua University, Beijing 100084, China

Abstract

Let $\chi'_d(n,q)$ (resp. $\chi_d(n,q)$) denote the minimum number of colors necessary to color a q-ary n-cube so that no two vertices that are at a distance at most d (resp. exactly d) get the same color. These two problems were proposed in the study of scalability of optical networks. In this paper, we provide upper and lower bounds on $\chi'_d(n,q)$ and $\chi_d(n,q)$ when q is a prime power.

Keywords: q-ary n-cube, Coloring problem, Coding theory, Vertex coloring

1 Introduction

Let $q = p^m$, where p is a prime. Let V_n be the n-dimensional vector space over a finite field $\mathbb{F}_q$, i.e.,

$$V_n = \{(x_1, \ldots, x_n) | x_i \in \mathbb{F}_q\}.$$

For $\mathbf{x} = (x_1, \ldots, x_n) \in V_n$ and $\mathbf{y} = (y_1, \ldots, y_n) \in V_n$, the Hamming distance $d_H(\mathbf{x}, \mathbf{y})$ between $\mathbf{x}$ and $\mathbf{y}$ is the number of the coordinates in which they differ, i.e.,

$$d_H(\mathbf{x}, \mathbf{y}) = |\{i | x_i \neq y_i,\ 1 \leq i \leq n\}|.$$

The Hamming weight $w_H(\mathbf{x})$ of a vector $\mathbf{x} \in V_n$ is the number of nonzero coordinates in $\mathbf{x}$. Obviously,

$$d_H(\mathbf{x}, \mathbf{y}) = w_H(\mathbf{x} - \mathbf{y}).$$

Let $G = (V, E)$ be a simple undirected graph with the vertex set V and the edge set E. For two distinct vertices $\mathbf{u}$ and $\mathbf{v}$ in V, the distance between $\mathbf{u}$ and $\mathbf{v}$, denoted by $d(\mathbf{u}, \mathbf{v})$, is the number of edges in a shortest path joining $\mathbf{u}$ and $\mathbf{v}$ and the diameter of G is the maximum distance between any two vertices of G.

A *q-ary n-cube* [1], denoted by $Q_n(q)$, is an undirected graph with the vertex set V_n and the edge set

$$E_n = \{(\mathbf{x}, \mathbf{y}) | \mathbf{x}, \mathbf{y} \in V_n, d_H(\mathbf{x}, \mathbf{y}) = 1\}.$$

The 2-ary n-cube $Q_n(2)$ is a hypercube suggested by Sullivan and Bashkow [14] and is one of the most popular, versatile and efficient topological structures of interconnection networks. By the definition, $Q_n(q)$ is $n(q-1)$-regular of order q^n, $d(\mathbf{x}, \mathbf{y}) = d_H(\mathbf{x}, \mathbf{y})$ for any distinct vertices $\mathbf{x}, \mathbf{y} \in V_n$ and the diameter of $Q_n(q)$ is n. Note that $Q_n(2)$ is a bipartite, but $Q_n(q)$ is not bipartite when $q \geq 3$.

*email: hanzhe101@gmail.com.
†email: mlu@math.tsinghua.edu.cn.

A coloring of V_n with L colors is a map Γ from the vertex set V_n to $\mathcal{L} = \{1, 2, \ldots, L\}$. A d-distance (resp. exactly d-distance) coloring of V_n is to color the vertices of V_n such that any two vertices with Hamming distance at most d (resp. exactly d) have different colors. Note that for a coloring of V_n with L colors

$$\Gamma : V_n \longrightarrow \mathcal{L},$$

it is a d-distance coloring of V_n if and only if for any two distinct vertices $\mathbf{x}, \mathbf{y} \in V_n$,

$$\Gamma(\mathbf{x}) \neq \Gamma(\mathbf{y}) \text{ if } d_H(\mathbf{x}, \mathbf{y}) \leq d$$

and it is an exactly d-distance coloring of V_n if and only if for any two distinct vertices $\mathbf{x}, \mathbf{y} \in V_n$,

$$\Gamma(\mathbf{x}) \neq \Gamma(\mathbf{y}) \text{ if } d_H(\mathbf{x}, \mathbf{y}) = d.$$

Denote $\chi'_d(n, q)$ (resp. $\chi_d(n, q)$) as the minimum number of colors needed for a d-distance (resp. an exactly d-distance) coloring of V_n. Clearly, $\chi_d(n, q) \leq \chi'_d(n, q)$ for each n.

These two coloring problems originally arose in the study of the scalability of optical networks [12]. Some bounds on $\chi'_d(n, 2)$ and $\chi_d(n, 2)$ were given and some exact values were determined (see for example [3]-[11], [13]-[15]). In the paper, we will give some lower and upper bounds on $\chi_d(n, q)$ and $\chi'_d(n, q)$. In particular, some bounds are given by using methods in coding theory.

The d-distance coloring and exactly d-distance coloring of V_n are equivalent to certain partitions of V_n, which are related to codes in coding theory. So we first introduce some definitions in coding theory. A nonempty subset C of V_n is called a q-ary code of length n. Any element in C is called a codeword of C. The minimum distance $d(C)$ is defined as the minimum Hamming distance between two distinct codewords of C. A q-ary code C of length n and minimum distance d is called an $(n, d)_q$ code. A q-ary code C of length n and minimum distance at least d is called an $(n, \geq d)_q$ code. Let $A_q(n, d)$ (resp. $A_q(n, \geq d)$) denote the maximum size of an $(n, d)_q$ (resp. $(n, \geq d)_q$) code. A q-ary code C is called a q-ary $[n, k]_q$ linear code if C is a k-dimensional subspace of V_n. A q-ary $[n, k]_q$ linear code with minimum distance d is called a q-ary $[n, k, d]_q$. For the fixed n and minimum distance d, let $k(n, d)_q$ denote the maximum dimension of a q-ary $[n, k, d]_q$ code. A q-ary code C of length n is called an $(n, \overline{\{d\}})_q$ forbidden distance code if $d_H(\mathbf{u}, \mathbf{v}) \neq d$ for any two distinct codewords $\mathbf{u}, \mathbf{v} \in C$. Given n and d, let $Q(n, d)_q$ denote the maximum size of a q-ary $(n, \overline{\{d\}})_q$ forbidden distance code.

An $(n, L, d)_q$-partition (resp. $(n, L, \overline{\{d\}})_q$-partition) of V_n is a set of subsets $\{B_i\}_{i=1}^L$ of V_n satisfying (i) $B_i \bigcap B_j = \emptyset$ for $i \neq j$ and $\bigcup_{i=1}^L B_i = V_n$; (ii) each B_i is a q-ary $(n, \geq d)_q$ code (resp. $(n, \overline{\{d\}})_q$ forbidden distance code). It is well known that a d-distance coloring (resp. an exactly d-distance) of V_n with L colors is equivalent to an $(n, L, d+1)_q$-partition (resp. $(n, L, \overline{\{d\}})_q$-partition) of V_n. Hence $\chi'_d(n, q)$ (resp. $\chi_d(n, q)$) is the minimum number L of subsets in any $(n, L, d+1)_q$ (resp. $(n, L, \overline{\{d\}})_q$-partition) of V_n.

Since $q^n = |V_n| = \sum_{i=1}^L |B_i| \leq LA_q(n, \geq d+1) \leq LA_q(n, d+1)$ and $A_q(n, d)$ is decreasing in d, we have

$$\chi'_d(n, q) \geq \frac{q^n}{A_q(n, d+1)}. \tag{1}$$

Note that for a q-ary $[n,k,d+1]_q$ linear code C, the cosets of C form an $(n,q^{n-k},d+1)_q$-partition of V_n. Hence, if there exists a q-ary $[n,k,d+1]_q$ linear code, then

$$\chi'_d(n,q) \le q^{n-k}. \tag{2}$$

In particular,

$$\chi'_d(n) \le q^{n-k(n,d+1)}. \tag{3}$$

Furthermore, if $A_q(n,d+1) = q^{k(n,d+1)}$, i.e., $A_q(n,d+1)$ is attained by a q-ary $[n,k,d+1]_q$ linear code, then

$$\chi'_d(n,q) = q^{n-k(n,d+1)}. \tag{4}$$

Since q-ary Hamming code $[\frac{q^r-1}{q-1}, \frac{q^r-1}{q-1}-r, 3]_q$ and q-ary simplex code $[\frac{q^r-1}{q-1}, r, q^{r-1}]_q$ are optimal code, by (4) we have that

$$\chi'_2\left(\frac{q^r-1}{q-1}, q\right) = q^r, \tag{5}$$

and

$$\chi'_{q^{r-1}-1}\left(\frac{q^r-1}{q-1}, q\right) = q^{\frac{q^r-1}{q-1}-r}. \tag{6}$$

In the next two Sections, we give some lower and upper bounds on $\chi_d(n,q)$ and $\chi'_d(n,q)$ for general d.

2 Bounds of $\chi'_d(n,q)$

First, we present the lower bound of $\chi'_d(n,q)$.

Theorem 2.1 *$\chi'_d(n,q) \ge \sum_{i=0}^{d/2}(q-1)^i\binom{n}{i}$ if d is even and $\chi'_d(n,q) \ge q\sum_{i=0}^{(d-1)/2}(q-1)^i\binom{n-1}{i}$ if d is odd.*

Proof. Suppose d is even. We consider all vertices within distance $d/2$ from the vertex $(0,0,\cdots,0)$. The total number of these vertices is $\sum_{i=0}^{d/2}(q-1)^i\binom{n}{i}$. The distance between any two of them is at most d. Thus, these vertices should have distinct colors.

Now we suppose d is odd. we consider all vertices within distance $(d-1)/2$ of the vertex $(a,0,\cdots,0)$ for any $a \in \mathbb{F}_q$. The total number of these vertices is $q\sum_{i=0}^{(d-1)/2}(q-1)^i\binom{n-1}{i}$. The distance between any two of them is at most d. Thus, these vertices should have distinct colors. ∎

Note. By Theorem 2.1, $\chi'_d(n,2) \ge \sum_{i=0}^{d/2}\binom{n}{i}$ for even d and $\chi'_d(n,2) \ge 2\sum_{i=0}^{(d-1)/2}\binom{n-1}{i}$ for odd k, which had been shown by Kim et al in [7]. In addition, Theorem 2.1 can be proved by Singleton bound in coding theory easily.

Next, we give some upper bounds of $\chi'_d(n,q)$.

Theorem 2.2 *For integers n and d, we have*

$$\chi'_d(n,q) \le q^{\left\lfloor \log_q\left(\sum_{j=0}^{d-1}(q-1)^j\binom{n-1}{j}\right)\right\rfloor+1}. \tag{7}$$

Proof. First, we show that there exists a $t \times n$ matrix H with the property that any d columns of H are linearly independent over $\mathbb{F}_q$ and there exist $d+1$ dependent columns. This H is actually the parity check matrix of an $[n, n-t, d+1]_q$-code.

Now, we describe a procedure for constructing a $t \times n$ parity check matrix H by choosing its column vectors sequentially. The first column vector can be any nonzero vector. Suppose that we already have a set V of i vectors so that any d of them are linearly independent. The $(i+1)$th vector can be chosen as long as it is not the linear combination of any $d-1$ vectors in V. In other words, since we are working over the field $\mathbb{F}_q$, the total number of unallowable vectors is at most $(q-1)\binom{i}{1}+(q-1)^2\binom{i}{2}+\ldots+(q-1)^{d-1}\binom{i}{d-1}$. Consequently, as long as $(q-1)\binom{i}{1}+(q-1)^2\binom{i}{2}+\ldots+(q-1)^{d-1}\binom{i}{d-1} < q^t-1$, we can still add a new column to H. Therefore a $t \times n$ parity check matrix H can be constructed. It follows that $t = \left\lfloor \log_q \left(\sum_{j=0}^{d-1} (q-1)^j \binom{n-1}{j} \right) \right\rfloor + 1$ and so $\chi'_d(n,q) \le q^t$ by (2). ∎

Now we consider the bounds of $\chi'_d(n,q)$ for $d=2$.

For a positive integer k and a prime l, a k-dimensional l-adic representation of a nonnegative integer i is a k-dimensional vector $(x_1, x_2, \cdots, x_k)$, where $x_i \in \{0, 1, \ldots, l-1\}$, such that

$$x_1 \cdot q^{k-1} + x_2 \cdot q^{k-2} + \cdots + x_{k-1} \cdot q + x_k = i.$$

Theorem 2.3 *$(q-1)n+1 \le \chi'_2(n,q)$ and $\chi'_2(n,q) \le q^{\lceil \log_q n \rceil + 1}$ if $q = p$.*

Proof. Consider vertices $(0,0,\cdots,0)$, $(a,0,\cdots,0)$, $(0,a,\cdots,0)$, $\cdots$, and $(0,\cdots,0,a)$, where $a \in \mathbb{F}_q \setminus \{0\}$. Then the distance between any two vertices is at most 2. Therefore, they should have distinct colors. That is, $\chi'_2(n,q) \ge (q-1)n+1$.

Let $m = \lceil \log_q n \rceil + 1$ and $M_q(n)$ be an $n \times m$ matrix whose $(i+1)$th row r_i $(0 \le i \le n-1)$ has the form (b_i, n_i), where b_i is the $(m-1)$-dimensional q-adic representation $(x_{i1}, x_{i2}, \cdots, x_{i,m-1})$ of integer i and $n_i = \sum_{j=1}^{m-1} x_{ij} + 1 (\mod q)$. For example,

$$M_3(5) = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 2 \\ 0 & 2 & 0 \\ 1 & 0 & 2 \\ 1 & 1 & 0 \end{pmatrix}.$$

Denote colors by m-dimensional vectors over $\mathbb{F}_q$. For any vertex $\mathbf{v} = (v_1, v_2, \cdots, v_n)$, color $\mathbf{v}$ with the color $\mathbf{v}M_q(n)$, where all arithmetic operations are performed in $\mathbb{F}_q$. Then there are q^m colors in total. Let $\mathbf{u}$ and $\mathbf{v}$ be any two vertices with distance $d \le 2$, and use $r_i = (x_{i1}, x_{i2}, \cdots, x_{i,m-1}, n_i)$ to denote the $(i+1)$th row of $M_q(n)$, where $0 \le i \le n-1$. We consider two cases.

Case 1 $d = 1$.

Suppose $u_i \ne v_i$ for some $1 \le i \le n$. Then $\mathbf{u}M_q(n) - \mathbf{v}M_q(n) = (u_i - v_i)r_{i-1}$. By the definition of $M_q(n)$, $r_{i-1} \ne \mathbf{0}$. Thus $\mathbf{u}M_q(n) \ne \mathbf{v}M_q(n)$.

Case 2 $d = 2$.

Suppose $u_i \neq v_i$ and $u_j \neq v_j$ for some $1 \leq i < j \leq n$. Then

$$\mathbf{u}M_q(n) - \mathbf{v}M_q(n) = (u_i - v_i)r_{i-1} + (u_j - v_j)r_{j-1}.$$

Denote $u_i - v_i = a_i$ and $u_j - v_j = a_j$ for short. Note that $(x_{i1}, x_{i2}, \cdots, x_{i,m-1}) \neq (x_{j1}, x_{j2}, \cdots, x_{j,m-1})$ for $i \neq j$. If $a_i(x_{i1}, x_{i2}, \cdots, x_{i,m-1}) = a_j(x_{j1}, x_{j2}, \cdots, x_{j,m-1})$, then $a_i \neq a_j$ but $a_i \sum_{t=1}^{m-1} x_{it} = a_j \sum_{t=1}^{m-1} x_{jt}$. Since $n_i = \sum_{j=1}^{m-1} x_{ij} + 1 (\mod q)$, we have $a_i n_i \neq a_j n_j$, which implies $\mathbf{u}M_q(n) \neq \mathbf{v}M_q(n)$.

Therefore, if the distance of $\mathbf{u}$ and $\mathbf{v}$ is at most 2, then $\mathbf{u}M_q(n) \neq \mathbf{v}M_q(n)$, and so vertices $\mathbf{u}$ and $\mathbf{v}$ have different colors. Hence $\chi_2'(n,q) \leq q^{\lceil \log_q n \rceil + 1}$. ∎

Note. In [15], Wan showed that $\chi_2'(n,2) \leq 2^{\lceil \log_2(n+1) \rceil}$. But the inequality $\chi_2'(n,q) \leq q^{\lceil \log_q(n+1) \rceil}$ does not always hold for q-ary n-cube. For example, consider the case $q = 3$ and $n = 2$. Then $3^{\lceil \log_3(2+1) \rceil} = 3$. But $\chi_2'(2,3) = 3^2 > 3$ since the distance of any two vertices in $Q_2(3)$ is at most 2 and so they should have distinct colors.

3 Bounds of $\chi_d(n,q)$

In this section, we consider the bounds of $\chi_d(n,q)$. Since the diameter of $Q_n(q)$ is n, $\chi_d(n,q) = 1$ for any $d \geq n+1$. So we can assume $d \leq n$.

Theorem 3.1 *$\chi_d(n,q) \geq q$. Particularly, $\chi_n(n,q) = q$ and $\chi_1(n,q) = q$.*

Proof. Consider vertices $(x_1, x_2, \cdots, x_n)$ with $x_1 = x_2 = \cdots = x_d = a$ for any $a \in \mathbb{F}_q$ and $x_{d+1} = \cdots = x_n = 0$. Then the distance between any two vertices is d. Therefore, they should have distinct colors. That is, $\chi_d(n,q) \geq q$.

For any $a \in \mathbb{F}_q$, denote $S_a = \{(x_1, x_2, \cdots, x_n) \ : \ x_1 = a, x_j \in \mathbb{F}_q \text{ for } 2 \leq j \leq n\}$. Then $V_n = \bigcup_{a \in \mathbb{F}_q} S_a$. For vertices $\mathbf{u}, \mathbf{v} \in S_a$ and some $a \in \mathbb{F}_q$, the distance between them is at most $n-1$ and they could have the same color. Thus $\chi_n(n,q) \leq q$, which implies $\chi_n(n,q) = q$.

We consider the case when $d = 1$. Let $\mathbb{F}_q = \{0, \alpha^0, \ldots, \alpha^{q-2}\}$ where α is a primitive element of $\mathbb{F}_q$ and $\mathcal{L} = \{0, 1, \ldots, q-1\}$. Put $\Gamma(0,0,\ldots,0) = 0$,

$$\Gamma(x_1, \ldots, x_n) = \sum_{x_k \neq 0, x_k = \alpha^{i_k}, 1 \leq k \leq n} i_k + 1 \mod q,$$

where $x_1, \ldots, x_n \in \mathbb{F}_q$. For any vertices $\mathbf{x}$ and $\mathbf{y}$ with distance 1, they have the different colors by the above definition. Thus $\chi_1(n,q) \leq q$, which implies $\chi_1(n,q) = q$. ∎

Note. When $d < n$ and $d > 1$, $\chi_d(n,q) = q$ does not always hold since we can give two counterexamples. When $d = 3, n = 5, q = 3$, we can find four vertices $(0,0,0,0,0)$,$(1,1,1,0,0)$, $(2,2,2,0,0)$, and $(2,0,1,2,0)$ whose distance between any two vertices is 3, which means that these four vertices would have 4 different colors. Likewise, when $d = 5, n = 7, q = 3$, there are four vertices $(0,0,0,0,0,0,0)$, $(1,0,1,0,1,1,1)$, $(0,0,2,2,2,2,1)$, and $(0,1,1,1,2,1,0)$ whose distance between any two vertices is 3, which means that these four vertices would have 4 different colors. But the conclusion holds for $q = 2$ since $Q_n(2)$ is bipartite.

Theorem 3.2 *For even d, $\chi_d(n,q) \geq \max\{q, \lfloor 2n/d \rfloor\}$.*

Proof. Consider vertices $(\underbrace{a,\cdots,a}_{d/2},0,\cdots,0)$, $(\underbrace{0,\cdots,0}_{d/2},\underbrace{a,\cdots,a}_{d/2},0,\cdots,0)$, $(\underbrace{0,\cdots,0}_{d},\underbrace{a,\cdots,a}_{d/2},0,\cdots,0)\cdots$, where $a \in \mathbb{F}_q \setminus \{0\}$. Then the distance between any two vertices is d. Therefore, they should have distinct colors. ∎

In the following we give some results about the exactly d-distance coloring of V_n by using the argument of linear forbidden distance codes. A q-ary $[n,k]_q$ linear code C is called a q-ary $[n,k,\overline{d}]_q$ linear forbidden distance code if C is also an $(n,\overline{d})$ forbidden distance code, i.e., $w_H(\mathbf{c}) \neq d$ for any nonzero codeword $\mathbf{c}$ of C. Given n and d, the maximum dimension of a q-ary $[n,k,\overline{d}]_q$ linear forbidden distance code is denoted by $k(n,\overline{d})$. Similar to d-distance coloring, we have

$$\chi_d(n,q) \leq q^{n-k}. \tag{8}$$

$$\chi_d(n,q) \leq q^{n-k(n,\overline{d})}. \tag{9}$$

In the following lemma the lower bound on $k(n,\overline{d})$ is given.

Lemma 3.3 *We have*

$$k(n,\overline{d}) \geq n - \left\lceil \log_q[2 + \binom{n-1}{d-1}(q-1)^{d-1}] \right\rceil. \tag{10}$$

Proof. First if

$$2 + \binom{n-1}{d-1}(q-1)^{d-1} \leq q^m, \tag{11}$$

then there exists a q-ary matrix H of size $m \times n$ such that any column of H is not equal to the $\mathbb{F}_q$-linear combination of any other $d-1$ columns of H. Hence there is a q-ary $[n,k,\overline{d}]_q$ linear forbidden distance code, where the dimension $k \geq n-m$. Take

$$m = \left\lceil \log_q[2 + \binom{n-1}{d-1}(q-1)^{d-1}] \right\rceil.$$

Thus,

$$\begin{aligned} k(n,\overline{d}) &\geq dim(C) \\ &\geq n-m \\ &\geq n - \left\lceil \log_q[2 + \binom{n-1}{d-1}(q-1)^{d-1}] \right\rceil. \end{aligned}$$

This completes the proof. ∎

Theorem 3.4 *We have*

$$\chi_d(n,q) \leq q^{\lceil \log_q[2+\binom{n-1}{d-1}(q-1)^{d-1}] \rceil}. \tag{12}$$

Proof. The theorem follows from Lemma 3.3 and (9). ∎

Acknowledgements

This work is partially supported by National Natural Science Foundation of China (Nos. 61373019, 11171097 and 61170289).

References

[1] L.N. Bhuyan and D.P. Agrawal, Generalized hypercube and hyperbus atructures for a cumputer network, IEEE Transactions on Computers, 33 (1984) 323-333.

[2] C. Bracken, Y.M. Chee and P. Purkayastha, Optimal family of q-ary codes obtained from a substructure of generalised Hadamad matrices, IEEE International Symposium on Information Theory Proceedings, 116-119, 2012.

[3] H. Enomoto, P. Frankl, N. Ito, K. Nomura, Codes with given distances, Graphs Combin. 3 (1987) 25-38.

[4] P. Frankl, Orthogonal vectors in the n-dimensional cube and codes with missing distances, Combinatorica 6 (1986) 279-285.

[5] F.-W. Fu, S. Ling and C. Xing, New results on two hypercube coloring problem, Discrete Applied Mathematics, 161(2013), 2937-2945.

[6] R.E. Jamison, G.L. Matthews, Distance k colorings of Hamming graphs, Congr. Numer. 183 (2006) 193-202.

[7] D.S. Kim, D.-Z. Du, P.M. Pardalos, A coloring problem on the n-cube, Discrete Appl. Math. 103 (2000) 307-311.

[8] W. Klotz, E. Sharifiyazdi, On the distance chromatic number of Hamming graphs, Adv. Appl. Discrete Math. 2 (2) (2008) 103-115.

[9] H.Q. Ngo, D.-Z. Du, R.L. Graham, New bounds on a hypercube coloring problem and linear codes, in: Proceedings of the International Conference on Information Technology: Coding and Computing, ITCC'01, IEEE, 2001, 542-546.

[10] H.Q. Ngo, D.-Z. Du and R.L. Graham, New bounds on a hypercube coloring problem, Inform. Process. Lett. 84 (2002) 265-269.

[11] P.R.J. Östergård, On a hypercube coloring problem, J. Combin. Theory Ser. A 108 (2004) 199-204.

[12] A. Pavan, P.-J. Wan, S.-R. Tong and D.H. Du, A new multihop lightwave network based on the generalized de-Bruijn graph, in: Proc. on IEEE INFOCOM, 1996, 498-507.

[13] Z. Skupień, BCH codes and distance multi- or fractional colorings in hypercubes asymptotically, Discrete Math. 307 (2007) 990-1000.

[14] H. Sullivan and T. R. Bashkow, A large scale homogeneous full distributed parallel machine, I. Proceeding of 4th Annual Symposium on Computer Architecture, 1977, 105-117.

[15] P.-J. Wan, Near-optimal conflict-free channel set assignments for an optical cluster-based hypercube network, J. Comb. Optim. 1 (1997) 179-186.